For Lateral Entry (Pharmacy)

- 1. Morphine is present in
 - (A) Atropa belladonna (B) Papaver somniferum
 - (C) Ricinus communis (D) Solanum nigrum
- 2. Gray baby syndrome is due to the indiscriminate use of
 - (A) Streptomycin (B) Chloramphenicol (C) Penicillin (D) Tetracycline
- 3. Which of the following is not a neurotransmitter ?
 - (A) Heparin (B) Nor adrenaline (C) Hydroxytryptamine (D) Prostaglandin
- 4. Wagner's test is used to detect the presence of

(A) Steroids (B) Alkaloids (C) Glycosides (D) Terpenes

- 5. One nanometer (nm) is equal to
 - (A) 10^{10} cm (B) 10^{12} cm (C) 10^{-7} cm (D) 10^{-4} cm
- 6. Indian and African Senna leaves differ from each other with respect to(A) Vein islet number (B) Stomatal index (C) Colour (D) All of the above
- 7. The conc. of sucrose in Simple Syrup BP is
 (A) 85% w/w (B) 60.7% w/w (C) 66.7% w/w (D) 66.7% w/v
- 8. The HLB value of Sodium lauryl sulphate is
 - (A) 4.0 (B) 4.7 (C) 7.4 (D) 40.0
- 9. The disintegration time for enteric coated tablet is
 - (A) 15 minutes (B) 30 minutes (C) 105 minutes (D) 135 minutes
- 10. Antibiotic which interacts with calcium ion is

(A) Erythromycin (B) Streptomycin (C) Tetracyclin (D) Ampicillin

- 11. Flow rate of granules from the hopper can be improved by adding(A) Lubricant (B) Disintegrant (C) Binder (D) Glidant
- 12. Sterilization temperature for aqueous solution in autoclave is(A) 60°c (B) 121°c (C) 160°c (D) 180°c
- 13. The vitamin administered with isoniazid to minimize its adverse reaction is(A) Vitamin A (B) Pyridoxine (C) Biotin (D) Pantothenic acid
- 14. State Pharmacy council should have the following number of elected members
 - (A) Six (B) Nine (C) Five (D) Seven
- 15. Synthesis of Urea takes place exclusively

(A) Kidney (B) Liver (C) Gall bladder (D) Urinary bladder

- 16. How many parts of 10% alcohol be mixed with 2 parts of 15% alcohol to get 12% alcohol ?
 - (A) 2 (B) 3 (C) 5 (D) 6
- 17. The UV- Visible region in the electromagnetic spectrum of radiation is

(A) 200-400 nm (B) 300-600 nm (C) 400-800 nm (D) 200-800 nm

- In Drugs & Cosmetics Act and Rules there under, list of substances that should by retail on prescription of Registered Medical Practitioner is given in
 - (A) Schedule H (B) Schedule G (C) Schedule J (D) Schedule O
- 19. Alkaloids in cinchona bark are detected by
 - (A) Iodine test (B) Thalleioquin test
 - (C) Liebermann-Burchard test (D) Nessler's test

20. The sweetening agent commonly use	ed in chewable tablet formula is	
(A) Sucrose (B) Cyclamate sodium ((C) Saccharin sodium (D) Mannitol	
21. The pH of a Pharmaceutical buffer sy	stem can be calculated by	
(A) pH partition theory	(B) Noyes-Whitney law	
(C) Henderson-Hasselbatch equation	(D) Michaelis-Menten equation	
22. The Pharmacy Council of India is con	nstituted by the	
(A) Central Government	(B) State Government	
(C) Parliament	(D) Legislative assembly	
23. Sigma blade mixers are commonly us	sed in	
(A) Wet granulation	(B) Dry granulation	
(C) Powder mixing	(D) Crude fibre mixing	
24. Water for injection differs from steril	le distilled water as it is free from	
(A) carbon dioxide (B) pyrogens (C)) preservatives (D) antioxidant	
25. Which of the following steroids show action?	vs predominant mineralocorticoid	
(A) Hydrocortisone	(B) Spiranolactone	
(C) Dexamethasone	(D) Fludrocortisone	
26. The Shick test is used to determine susceptibility to		
(A) Measles (B) Diptheria (C) Pol	lio (D) Typhoid	
27. Disposable syringes are made up of		
(A) Polypropylene	(B) Transparent polystyrene	
(C) Glass	(D) Poly Tetra Chloro Ethylene	

- 28. Choose the correct pH of the lachrymal fluid
 - (A) 8.0 (B) 6.2 (C) 7.4 (D) 9.0
- 29. A drug which has antipyretic, anti-inflammatory and antiplatelet activity is(A) Sulfinpyrazone (B) Aspirin (C) Ticlopidine (D) Acetaminophen
- 30. Cocaine is used as a
 - (A) Topical anaesthetic (B) Cough suppressant
 - (C) Stimulant (D) Respiratory infections
- 31. Penicillinase resistant penicillin is
 - (A) Amoxycillin (B) Ampicillin (C) Penicillin V (D) Methicillin
- 32. Tyndallisation means
 - (A) Successive autoclaving with a bactericide
 - (B) Successive heating with a bactericide
 - (C) Successive heating at low temperature
 - (D) Successive heating at low temperature and incubation
- 33. Vincristine and vinblastine act by
 - (A) Binding with the protein tubulin and arrest at metaphase
 - (B) Inhibiting the protein synthesis
 - (C) Acting as antimetabolite
 - (D) Inhibiting the enzyme system
- 34. Sulphonamide tragedy was due to its combination with
 - (A) Penicillin (B)Streptomycin (C)Diethylene hydrochloride (D)Bicarbonate

35. (Cholinergic	receptor	present	on	intestinal	muscle	is
-------	-------------	----------	---------	----	------------	--------	----

(A) Histamine receptor		(B) Muscarin	ic receptor
(C) Nicotine receptor	((D) Beta rece	ptor
36. Metronidazole inhibits anaerobic bacteria and protozoa by			otozoa by
(A) Affecting the structure	re of DNA	molecule of	the organism
(B) Destroying the riboso	omes		
(C) Inhibiting the cytochrome system			
(D) Inhibiting the protein	synthesis		
37. Antihypertensive drug inhibits the renin-angiotensin system is			
(A) Reserpine (B) Capto	opril (C) M	Iethyl dopa (D) Propanolol
38. Titanium dioxide is commonly present in			
(A) Vanishing cream	(B) Sunscreen	cream
(C) Aqueous calamine cr	eam (E	D) Opthalmic	cream
39. Powdered digitalis is dried at a temperature			
(A) Not exceeding 60°c	(B) 65°c	(C) 75°c	(D) 100°c
40. The Mantoux test uses			

(A) Old tuberculin	(B) Diptheria toxin
--------------------	---------------------

(C) Serum antigen (D) Polysaccharide antigen

41. The neurotransmitter released at the end of sympathetic nerve fibre is

(A) Epinephrine (B) Nor-epinephrine (C) Acetylcholine (D) Physostigmine

42. Rancidity of a fat is due to

(A) Oxidation (B) Saponification (C) Hydrolysis (D) Neutralisation

43. The vitamin which has deodorant property is

(A) Vitamin A (B) Vitamin C (C) Vitamin D (D) Vitamin E

44. Erythromycin is an antibiotic. It belongs to the class of

(A) ß-lactum (B) Aminoglycosides (C) Macrolide (D) Peptide

- 45. The principal structural component of the cell wall in bacteria ia made up of
 - (A) simple protein (B) peptidoglycan polymer
 - (C) complex polysaccharides (D) glycoprotein
- 46. Tetracyclines are avoided during pregnancy because
 - (A) It is teratogenic (B) it may affect the bone growth of foetus
 - (C) it causes discolouration of mothers teeth (D) it may cause abortion
- 47. The tear secretion contains an antibacterial enzyme known as

(A) Zymase (B) Diastase (C) Lysozyme (D) Lipase

48. Schedule M as per Drugs and Cosmetics Act is concerned with

- (A) Opthalmic preparations (B) Life period of drugs
- (C) Requirements of factory premises (D) List of prescription drugs

49. Varicella zoster is the causative organism for

(A) Small pox (B) Dermatophytosis (C) Herpes (D) Tuberculosis

50. The statement 'store in a cold place' as per IP, means

(A) store at room temperature (B) store between 2° to 8° c

(C) store between 8° to 25° c (D) store at 0° c

51. Durability of a tablet to combined effects of shock and abrasion is evaluated by using

(A) Hardness tester	(B) Disintegration test apparatus	
(C) Friabilator	(D) Screw Gauge	
52. Emetine is the predominant alkaloid of		
(A) Rauwolfia serpentina	(B) Vinca rosea	
(C) Cephaelis ipecacuanha	(D) Swertia chirata	
53. The entire plant of Swertia chirata is used as		
(A) Cardiotonic (B) Bitter tonic (C) Diuretic (D) Anti diabetic		
54. Clark's formula for calculation of dose of a drug is based on		
(A) Age (B) Body weight (C) Surface area (D) Drug activity		
55. The organism that enter the body only through an injury is		
(A) Salmonella typhi	(B) Clostridium tetani	
(C) Clostridium botulinum	(D) Streptococcus pyrogenes	
56. The pH range of methyl orange is		
(A) 2.9-4.6 (B) 1.9-3.6	(C) 3.9- 5.6 (D) 4.2-6.3	
57. In the preparation of ointments, macrogols are used as		
(A) Water soluble base	(B) Hydrocarbon base	
(C) Absorption base	(D) Oleogenous base	

58. The particle size of the dispersed solids in a suspension is usually greater than

(A) $0.5 \ \mu m$ (B) $1 \ \mu m$ (C) $5 \ \mu m$ (D) $10 \ \mu m$

- 59. In the extemporaneous preparation of a suspension, Levigation is used to
 - (A) Reduce the zeta potencial (B) avoid bacterial growth
 - (C) reduce particle size (D) enhance viscosity

60. In order to produce characteristic pharmacological action, a drug must always

- (A) reach high blood levels (B) absorbed from GIT readily
- (C) achieve adequate conc. at site of action (D) excrete unchanged in urine

Answers:

1. B	21. C	41. B
2. В	22. A	42. A
3. A	23. C	43. B
4. B	24. B	44. C
5. C	25. D	45. B
6. D	26. B	46. C
7. C	27. В	47. C
8. D	28. C	48. C
9. D	29. B	49. C
10. C	30. A	50. B
11. D	31. D	51. C
12. B	32. C	52. C
13. B	33. A	53. B
14. D	34. A	54. B
15. A	35. B	55. B
16. B	36. A	56. A
17. D	37. В	57. A
18. A	38. B	58. A
19. B	39. A	59. C
20. D	40. A	60. C